

DIFFERENCE

Contact and systemic fungicide against scab in apples and pears.

Emulsifiable concentrate formulation containing 250 g/l (23.6 % w/w) difenoconazole

MAPP No.: 16129

GROUP

3

FUNGICIDE

FUNGICIDE

Contents:

1Le



Production date / Batch number:
see packaging

FOR USE ONLY AS A
HORTICULTURAL FUNGICIDE

FOR PROFESSIONAL USE ONLY

Important : The (COSHH) Control
of Substances Hazardous to Health
Regulations may apply to the use
of this product at work.

Marketing company:

Q-Chem nv
Leeuwerweg 138
B-3803 Sint Truiden - Belgium
Tel: +32 (0)11 78 57 17
Fax: +32 (0)11 68 15 65 A

AUTHORISATION HOLDER:

 **GLOBACHEM NV**
Lichtenberglaan 2019
BE-3800 Sint-Truiden
www.globachem.com

@ registered trademark of
Globachem NV

Distributed by: **Certis Belchim BV**
Suite 5, 3 Riverside | Granta Park | Great Abington | Cambridgeshire | CB21 6AD
Tel.: 01223 652500 | info.uk@certisbelchim.com | www.certisbelchim.com

DIFFERENCE - Emulsifiable concentrate containing 250 g/l (23.6 % w/w) difenoconazole

H302 Harmful if swallowed
H304 May be fatal if swallowed and enters airways
H319 Causes serious eye irritation
H411 Toxic to aquatic life with long lasting effects


P102 Keep out of reach of children
P280 Wear eye protection
P301+P310 IF SWALLOWED: immediately call a POISON
CENTRE or doctor/physician

P331 Do NOT induce vomiting
P337+P313 If eye irritation persists: Get medical advice/
attention

P501 Dispose of contents/container to a licensed hazardous-waste
disposal contractor or collection site except for empty clean
containers which can be disposed of as non-hazardous waste

**EUH401 To avoid risks to human health and the environment, comply with
the instructions for use**

READ THE PRECAUTIONS ON THE BOTTLE LABEL BEFORE USE



IMPORTANT INFORMATION - FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Crops	Max. individual dose (L/ha)	Max number of applications (per crop)	Latest time of application
Apples, pears	0.2	4	14 days before harvest

Other specific restrictions:

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme for broadcast air-assisted sprayers. Before each spraying operation from a broadcast air-assisted sprayer, either a LERAP must be carried out in accordance with the 'Local Environment Risk Assessment for Pesticides Broadcast Air-Assisted Sprayers' booklet available from the HSE Chemicals Regulation Directorate's website or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years. Do not re-use container for any purpose.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH
THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION
PRODUCTS**

For warehouse
purpose only:



Certis Belchim
GROWING TOGETHER

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practical in addition to the following personal protective equipment:

* WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

* WASH HANDS AND EXPOSED SKIN before meals and after work.

* WHEN USING, DO NOT DRINK, EAT OR SMOKE.

If you feel unwell, seek medical advice immediately (show this container or label).

* IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical attention.

Environmental Protection

* DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

* DO NOT ALLOW DIRECT SPRAY from broadcast air-assisted sprayers to fall within 20 m of the top of the bank of a static or flowing waterbody unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 5 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

Storage and Disposal

* KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

* WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

* KEEP OUT OF REACH OF CHILDREN.

* KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

* Use appropriate containment to avoid environmental contamination.

* Do not re-use container for any purpose.

* Store in a cool, dry place.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CROP SPECIFIC INFORMATION

APPLES AND PEARS

DIFFERENCE controls apple scab (*Venturia inaequalis*) and pear scab (*Venturia pirina*) in apples and pears. It can be applied from bud burst to 14 days before harvest (including during flowering) up to four times in a season with a minimum interval of 8 days.

Dose: 0.2 L/ha.

Alternate products with different modes of action in your spray programme against scab (see FRAC fungicide group list on the FRAC website).

CONSULT PROCESSOR BEFORE USING ON CROPS FOR PROCESSING.

MIXING AND SPRAYING

Mixing

Make sure the sprayer is set to give an even application at the correct volume.

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of DIFFERENCE and continue agitation while adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying.

Thoroughly wash all spray equipment (tank and nozzles) at least 2 times with water and add a cleaning agent after use.

Spray volume

Apples and pears: 200-1000 litres water per hectare.

Spraying

Apply as a MEDIUM quality spray as defined by BCPC. A spray pressure of 2-3 bars is preferred.

COMPATIBILITY

In case of tank mixes with trace elements, it is recommended that DIFFERENCE is added first to the tank with constant agitation and the mixture is sprayed without delay.

Always use constant agitation of the sprayer tank during mixing, transportation and application. Spray immediately.

RESISTANCE

DIFFERENCE contains a DMI fungicide. Resistance to some DMI fungicides has been identified in scab (*Venturia inaequalis* and *Venturia pirina*) which may seriously affect the performance of some products. For some further advice on resistance management in DMI's contact your agronomist or specialist advisor, and visit the FRAG-UK website.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



WEBLINK to specific data sheet on manufacturer's website:

www.certisbelchim.co.uk/pdf/MSDS/Difference_MSDS.pdf

Alternatively, contact your supplier.