

DIFFERENCE®

MAPP 16129

GROUP 3 FUNGICIDE

Contact and systemic fungicide against scab in apples and pears.

Emulsifiable concentrate formulation containing 250 g/l (23.6 % w/w) difenoconazole.

FUNGICIDE

F

Net content

1Le

FOR PROFESSIONAL USE ONLY

Production date / Batch number:
see packaging

For advice on medical emergencies, fires, spillages or chemical hazards ONLY – phone Carechem 24 EU: 01235 239670

For technical and non-emergency calls phone 0845 373 0305

Marketing Company
Q-Chem NV
Leeuwerweg 138
B-3803 Sint Truiden - Belgium
Tel: +32 11 78 57 17


Globachem
Your Crop Counts

Authorisation Holder
GLOBACHEM NV
Lichtenberglaan 2019
BE-3800 Sint-Truiden
www.globachem.com
Tel: +32 11 78 57 17

© Registered Trademark of Globachem NV

IMPORTANT INFORMATION

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Crops	Maximum individual dose	Maximum number of applications	Latest time of application	Other specific restrictions
Apples, pears	Refer to leaflet attached to the product pack for full details.			

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

THE CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH (COSHH) REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK.

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practical in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection. WASH HANDS AND EXPOSED SKIN before meals and after work. WHEN USING, DO NOT DRINK, EAT OR SMOKE.

If you feel unwell, seek medical advice immediately (show this container or label). IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical attention.

Environmental Protection

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. DO NOT ALLOW DIRECT SPRAY from broadcast air-assisted sprayers to fall within 20 m of the top of the bank of a static or flowing waterbody unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 5 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely. KEEP OUT OF REACH OF CHILDREN. KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. Use appropriate containment to avoid environmental contamination. Do not re-use container for any purpose. Store in a cool, dry place.

DIFFERENCE®

Emulsifiable concentrate containing 250 g/l (23.6% w/w) difenoconazole.

UFI: WC00-509Q-400N-GX5A

DANGER

H302 Harmful if swallowed.

H304 May be fatal if swallowed and enters airways.

H319 Causes serious eye irritation.

H351 Suspected of causing cancer (NI-only).

H410 Very toxic to aquatic life with long lasting effects.

P102 Keep out of reach of children.

P280 Wear protective gloves/ protective clothing/ eye protection/ face protection.

P301 + P310 IF SWALLOWED: Immediately call a POISON CENTRE or doctor.

P331 Do NOT induce vomiting.

P305+P351+P338 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

P337+P313 If eye irritation persists get medical advice/attention.

P391 Collect spillage.

P501 Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean, triple rinsed containers which can be disposed of as non-hazardous waste. EUH066 Repeated exposure may cause skin dryness or cracking.

EUH401 To avoid risks to human health and the environment, comply with the instructions for use.

05250101_2532 • 107044171-UK



Certis Belchim
GROWING TOGETHER

Distributed by: **Certis Belchim BV**
Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridgeshire CB21 6AD
Tel: 01223 652500 | Email: info.uk@certisbelchim.com | www.certisbelchim.co.uk

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS A HORTICULTURAL FUNGICIDE

Crops	Maximum individual dose (L/ha)	Maximum number of applications (per crop)	Latest time of application
Apples, pears	0.2	4	14 days before harvest

Other specific restrictions

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme for broadcast air-assisted sprayers. Before each spraying operation from a broadcast air-assisted sprayer, either a LERAP must be carried out in accordance with the 'Local Environment Risk Assessment for Pesticides Broadcast Air-Assisted Sprayers' booklet available from the HSE Chemicals Regulation Directorate's website or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years. Do not re-use container for any purpose.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CROP SPECIFIC INFORMATION

APPLES AND PEARS

DIFFERENCE® controls apple scab (*Venturia inaequalis*) and pear scab (*Venturia pirina*) in apples and pears. It can be applied from bud burst to 14 days before harvest (including during flowering) up to four times in a season with a minimum interval of 8 days.

Dose: 0.2 L/ha.

Alternate products with different modes of action in your spray programme against scab (see FRAC fungicide group list on the FRAC website).

CONSULT PROCESSOR BEFORE USING ON CROPS FOR PROCESSING.

MIXING AND SPRAYING

Mixing

Make sure the sprayer is set to give an even application at the correct volume.

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of DIFFERENCE® and continue agitation while adding the rest of the water.

Agitate the mixture thoroughly before use and continue agitation during spraying.

Thoroughly wash all spray equipment (tank and nozzles) at least 2 times with water and add a cleaning agent after use.

Spray volume

Apples and pears: 200-1000 litres water per hectare.

Spraying

Apply as a MEDIUM quality spray as defined by BCPC. A spray pressure of 2-3 bar is preferred.

COMPATIBILITY

In case of tank-mixes with trace elements, it is recommended that DIFFERENCE® is added first to the tank with constant agitation and the mixture is sprayed without delay.

Always use constant agitation of the sprayer tank during mixing, transportation and application. Spray immediately.

RESISTANCE

DIFFERENCE® contains a DMI fungicide. Resistance to some DMI fungicides has been identified in scab (*Venturia inaequalis* and *Venturia pirina*) which may seriously affect the performance of some products. For some further advice on resistance management in DMI's contact your agronomist or specialist advisor, and visit the FRAG-UK website.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:

WEBLINK to specific data sheet on manufacturer's website:

https://certisbelchim.co.uk/pdf/MSDS/Difference_MSDS.pdf

Alternatively, contact your supplier.

